



Uniform Data System 2015-2016 Clinical Quality Measures Crosswalk

Clinical Focus Area 2016	Performance Measure 2016	Numerator Description 2016	Denominator Description 2016	Major Differences in Criteria between 2016 and 2015
Body Mass Index (BMI) Screening and Follow-Up Plan	<p>Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous 6 months AND when the BMI is outside of normal parameters, for whom a follow-up plan is documented during the encounter or during the previous 6 months of the current encounter</p> <p>Normal parameters:</p> <ul style="list-style-type: none"> • Age 18 - 64 years BMI => 18.5 and < 25 kg/m² • Age 65 years and older BMI => 23 and < 30 kg/m² 	Patients with a documented BMI (not just height and weight) during their most recent visit or during the previous six months of the most recent visit, and when the BMI is outside of normal parameters, a follow-up plan is documented during the visit or during the previous six months of the current visit	All patients 18 years of age or older during the measurement year, who had at least 1 medical visit during the reporting year	<p>Additional patient exclusions:</p> <ul style="list-style-type: none"> • receiving palliative care, • refusing measurement of height and/or weight, • in an urgent or emergent medical situation, or • if there are other reasons documented in the medical record
Cervical Cancer Screening	Percentage of women 21-64 years of age, who received 1 or more Pap tests to screen for cervical cancer	Women with 1 or more Pap tests during the measurement period or the 2 years prior to the measurement period	Women 23-64 years of age with a visit during the measurement period	<ul style="list-style-type: none"> • Age criteria changed to age 23 through 64 years (previously age 24 through 64 years) • No longer includes concurrent human papillomavirus and Pap test for those age 30 and older

Clinical Focus Area 2016	Performance Measure 2016	Numerator Description 2016	Denominator Description 2016	Major Differences in Criteria between 2016 and 2015
Childhood Immunization Status	Percentage of children 2 years of age who had 4 diphtheria, tetanus, and acellular pertussis (DTaP); three polio (IPV), 1 measles, mumps, and rubella (MMR); 3 H. influenza type B (HiB); 3 hepatitis B (Hep B); 1 chicken pox (VZV); 4 pneumococcal conjugate (PCV); 1 hepatitis A (Hep A); 2 or 3 rotavirus (RV); and 2 influenza (flu) vaccines by their second birthday	Children who have evidence showing they received recommended vaccines, had documented history of the illness, had a seropositive test result, or had an allergic reaction to the vaccine by their second birthday	Children who turn 2 years of age during the measurement period and who have a visit during the measurement period	<ul style="list-style-type: none"> • Age criteria changed to age 2 years (previously age 3 years) • Adds 1 Hep A, 2 or 3 RV, and 2 flu vaccines
Colorectal Cancer Screening	Percentage of adults 50-75 years of age who had appropriate screening for colorectal cancer	<p>Patients with 1 or more screenings for colorectal cancer</p> <p>Appropriate screenings are defined by any 1 of the following criteria:</p> <ul style="list-style-type: none"> • fecal occult blood test (FOBT) during the measurement period, • flexible sigmoidoscopy during the measurement period or the 4 years prior to the measurement period, or • colonoscopy during the measurement period or the 9 years prior to the measurement period 	Patients 50-75 years of age with a visit during the measurement period	Age 50 through 75 years (previously 51 through 74 years)

Clinical Focus Area 2016	Performance Measure 2016	Numerator Description 2016	Denominator Description 2016	Major Differences in Criteria between 2016 and 2015
Controlling High Blood Pressure	Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (< 140/90 mmHg) during the measurement period	Patients whose blood pressure at the most recent visit is adequately controlled (systolic blood pressure < 140 mmHg and diastolic blood pressure < 90 mmHg) during the measurement period	Patients 18-85 years of age who had a diagnosis of essential hypertension within the first 6 months of the measurement period or any time prior to the measurement period	<ul style="list-style-type: none"> Age 18 through 85 years (previously age 18 through 84 years) Excludes patients with dialysis or renal transplant before or during the measurement period and Stage 5 chronic kidney disease Two or more medical visits are no longer required
Coronary Artery Disease (CAD): Lipid Therapy	Percentage of patients aged 18 years and older with a diagnosis of CAD who were prescribed a lipid-lowering therapy	Number of patients who received a prescription for, or were provided, or were taking lipid lowering medications	Number of patients who were seen during the measurement year after their 18th birthday, who had at least 1 medical visit during the measurement year, at least 2 medical visits ever, and who had an active diagnosis of CAD including any diagnosis for myocardial infarction (MI) or who had had cardiac surgery in the past	N/A
Dental Sealants for Children aged 6 - 9 years	Percentage of children, age 6 through 9 years, at moderate to high risk for caries who received a sealant on a permanent first molar during the measurement period	Patients who received a sealant on a permanent first molar tooth in the measurement year	Dental patients aged 6 through 9 years who had an oral assessment or comprehensive or periodic oral evaluation visit during the measurement year and documented as having moderate to high risk for caries	Work provided by students will count as a visit (did not in 2015)
Diabetes: Hemoglobin A1c (HbA1c) Poor Control	Percentage of patients 18-75 years of age with diabetes who had HbA1c > 9.0% during the measurement period	Patients whose most recent HbA1c level is >9.0% (performed during the measurement period)	Patients 18-75 years of age with diabetes with a visit during the measurement period	<ul style="list-style-type: none"> Age 18 through 75 years (previously age 18 through 74 years) No longer has exclusions Two or more medical visits are no longer required
Early Entry into Prenatal Care	Percentage of prenatal care patients who entered treatment during their first trimester	Women entering prenatal care at the health center or with the referral provider during their first trimester	Women seen for prenatal care during the measurement period	N/A



Clinical Focus Area 2016	Performance Measure 2016	Numerator Description 2016	Denominator Description 2016	Major Differences in Criteria between 2016 and 2015
HIV Linkage to Care	Percentage of newly diagnosed HIV patients who had a medical visit for HIV care within 90 days of first-ever HIV diagnosis	Patients who had a medical visit for HIV care within 90 days of first-ever HIV diagnosis	Patients first diagnosed with HIV by the health center between October 1 of the prior year through September 30 of the current measurement year	N/A
Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antithrombotic	Percentage of patients 18 years of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG), or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had an active diagnosis of IVD during the measurement period, and who had documentation of use of aspirin or another antithrombotic during the measurement period	Patients who have documentation of use of aspirin or another antithrombotic during the measurement period	Patients 18 years of age and older with a visit during the measurement period, and an active diagnosis of IVD, or who were discharged alive for AMI, CABG, or PCI in the 12 months prior to the measurement period	N/A
Low Birth Weight	Percentage of babies born to health center patients whose birth weight was below normal (less than 2,500 grams)	Babies born with a birth weight of under 2,500 grams	Live births during the measurement period for women who received prenatal care from the health center or by a referral provider	N/A
Screening for Clinical Depression and Follow-Up Plan	Percentage of patients aged 12 years and older screened for clinical depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of the positive screen	Patients screened for clinical depression on the date of the encounter using an age appropriate standardized tool AND if positive, a follow-up plan is documented on the date of the positive screen	All patients aged 12 years and older before the beginning of the measurement period with at least one eligible encounter during the measurement period	Removes from denominator patients who: <ul style="list-style-type: none"> • refuse to participate, • urgent or emergent situations, or • if the patient's functional capacity or motivation to improve impacts the accuracy of results

Clinical Focus Area 2016	Performance Measure 2016	Numerator Description 2016	Denominator Description 2016	Major Differences in Criteria between 2016 and 2015
Tobacco Use Screening and Cessation Intervention	Percentage of patients aged 18 years and older who were screened for tobacco use 1 or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user	Patients who were screened for tobacco use at least once within 24 months AND who received tobacco cessation intervention if identified as a tobacco user	All patients aged 18 years and older seen for at least 2 visits or at least 1 preventive visit during the measurement period	<ul style="list-style-type: none"> • Adds documentation of medical reason(s) for not screening for tobacco use as an exception • Includes a broader set of visit types and services • Changes visits to seen at least twice or at least once for a preventive visit
Use of Appropriate Medications for Asthma	Percentage of patients 5-64 years of age who were identified as having persistent asthma and were appropriately prescribed medication during the measurement period	Patients who were dispensed at least 1 prescription for a preferred therapy during the measurement period	Patients 5-64 years of age with persistent asthma and a visit during the measurement period	<ul style="list-style-type: none"> • Age 5 through 64 years (previously age 5 through 40 years) • Removes criteria of 2 visits ever • Excludes patients with emphysema, chronic obstructive pulmonary disease (COPD), cystic fibrosis, or acute respiratory failure during or prior to the measurement period • Allergic reaction to asthma medications no longer an exclusion
Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents	Percentage of patients 3 - 17 years of age who had evidence of basal metabolic index (BMI) percentile documentation and who had documentation of counseling for nutrition and who had documentation of counseling for physical activity during the measurement period	Number of patients in the denominator who had their BMI percentile documented during the measurement year and who had documentation of counseling for nutrition and who had documentation of counseling for physical activity during the measurement period	Patients 3 through 17 years of age with at least one medical visit during the measurement period	N/A



Financial Focus Area 2016	Performance Measure 2016	Numerator Description 2016	Denominator Description 2016	Major Differences in Criteria between 2016 and 2015
BPHC Health Center Program Grant Cost Per Total Patient	Total BPHC section 330 grant funds per patient serviced in the measurement calendar year	Total accrued BPHC section 330 grants drawn-down for the period from January 1 to December 31, of the measurement calendar year	Total number of patients	N/A
Medical Cost Per Medical Patient	Total medical cost per medical visit in the measurement calendar year	Total accrued medical staff and other medical costs after allocation of overhead, excluding medical lab and x-ray costs	Medical visits, excluding nurse visits	N/A
Total Cost Per Total Patient	Total cost per patient serviced in the measurement calendar year	Total accrued cost before donations and after allocation of overhead	Total number of patients	N/A