

# 2017 Reporting Requirements

Focus Area (2016 e-CQMs reference)	Performance Measure	2017 Update
<b>Clinical Performance Measures</b>		
Diabetes: Hemoglobin A1c Poor Control ( <a href="#">CMS122v4</a> )	Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period	Revised to align with <a href="#">CMS122v5</a>  Changes to Measurement Guidance, Value Set, and Logic.
Controlling High Blood Pressure ( <a href="#">CMS165v4</a> )	Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (less than 140/90 mm Hg) during the measurement period	Revised to align with <a href="#">CMS165v5</a>  Changes to Guidance, Value Set, and Logic
Low Birth Weight	Percentage of babies of health center prenatal care patients born whose birth weight was below normal (less than 2,500 grams)	No Update
Early Entry into Prenatal Care	Percentage of prenatal care patients who entered prenatal care during their first trimester	No Update
Childhood Immunization Status (CIS) ( <a href="#">CMS117v4</a> )	Percentage of children 2 years of age who were fully immunized by their second birthday	Revised to align with <a href="#">CMS117v5</a>  Changes to Guidance, Value Set, and Logic.
Cervical Cancer Screening ( <a href="#">CMS124v4</a> )	Percentage of women 21-64 years of age, who were screened for cervical cancer using either of the following criteria: * Women age 21-64 who had cervical cytology performed every 3 years * Women age 30-64 who had cervical cytology/human papillomavirus (HPV) co-testing performed every 5 years	Revised to align with <a href="#">CMS124v5</a>  Reintroduces concurrent HPV and Pap test for those age 30-64. Changes to Value Set and Logic.
Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents ( <a href="#">CMS155v4</a> )	Percentage of patients 3 -17 years of age who had a medical visit and who had evidence of the following during the measurement period: height, weight, and BMI percentile documentation; and who had documentation of (1) counseling for nutrition; and (2) counseling for physical activity during the measurement year	Revised to align with <a href="#">CMS155v5</a>  Changes to Value Set and Logic.
Body Mass Index (BMI) Screening and Follow-Up ( <a href="#">CMS69v4</a> )	Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous six months AND when the BMI is outside of normal parameters, a	Revised to align with <a href="#">CMS69v5</a>

	follow-up plan is documented during the encounter or during the previous six months of the current encounter with the BMI outside of normal parameters: Age 18 years and older BMI => 18.5 and < 25 kg/m <sup>2</sup>	Changes to Initial Patient Population, Denominator Exclusions and Exceptions, Guidance, Value Set, and Logic.
Tobacco Use: Screening and Cessation Intervention ( <a href="#">CMS138v4</a> )	Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user	Revised to align with <a href="#">CMS138v5</a>  Changes to Guidance, Value Set, and Logic
Use of Appropriate Medications for Asthma ( <a href="#">CMS126v4</a> )	Percentage of patients 5-64 years of age with a diagnosis of persistent asthma and were appropriately ordered medication during the measurement period	Revised to align with <a href="#">CMS126v5</a>  Changes to Numerator, Value Set, and Logic
Coronary Artery Disease (CAD): Lipid Therapy	Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease (CAD) who were prescribed a lipid-lowering therapy	No Update
Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antithrombotic ( <a href="#">CMS164v4</a> )	Percentage of patients 18 years of age and older who were diagnosed with acute myocardial infarction (AMI), coronary artery bypass graft (CABG), or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had an active diagnosis of ischemic vascular disease (IVD) during the measurement period, and had documentation of use of aspirin or another antiplatelet during the measurement period	Revised to align with <a href="#">CMS164v5</a> Vascular Disease (IVD): Use of Aspirin or Another Antiplatelet  Changes from antithrombotic to antiplatelet, Changes to Title, Denominator, Initial Patient Population, Value Set, and Logic.
Colorectal Cancer Screening ( <a href="#">CMS130v4</a> )	Percentage of adults 50-75 years of age who had appropriate screening for colorectal cancer	Revised to align with <a href="#">CMS130v5</a> Changes to Value Set and Logic.
Screening for Clinical Depression and Follow-Up Plan ( <a href="#">CMS2v5</a> )	Percentage of patients aged 12 years and older screened for depression on the date of the visit using an age appropriate standardized depression screening tool AND if screening is positive, a follow-up plan is documented on the date of the positive screen	Revised to align with <a href="#">CMS2v6</a>  Clinical depression changed to depression. Changes to Logic and Value Set.
HIV Linkage To Care	Percentage of newly diagnosed HIV patients who were seen for follow-up treatment within 90 days of the first-ever HIV diagnosis	No Update
Dental Sealants for Children between 6 -9 Years ( <a href="#">CMS277v0</a> )	Percentage of children, age 6 through 9 years, at moderate to high risk for caries who received a sealant on a permanent first molar during the measurement period	No Update